

## A novel role of c-FLIP protein in regulation of ER stress response

Silvia Conti <sup>1</sup> - Simonetta Petrunaro <sup>1</sup> - Elettra Sara Marini <sup>2</sup> - Silvia Masciarelli <sup>1</sup> - Luana Tomaipitina <sup>1</sup> - Antonio Filippini <sup>1</sup> - [Claudia Giampietri](#) <sup>1</sup> - Elio Ziparo <sup>1</sup>

<sup>1</sup> Sapienza Università di Roma, Dipartimento di Scienze Anatomiche, Istologiche, Medico-Legali e dell'Apparato Locomotore, Roma, Italia - <sup>2</sup> University of Oslo, Department of Biosciences, Oslo, Norvegia

Cellular-Flice-like Inhibitory Protein (c-FLIP) is an apoptosis modulator known to inhibit the extrinsic apoptotic pathway thus blocking Caspase-8 processing in the Death Inducing Signalling Complex (DISC). We previously demonstrated that c-FLIP localizes at the Endoplasmic Reticulum (ER) and that c-FLIP-deficient Mouse Embryonic Fibroblasts (MEFs) display an enlarged ER morphology. In the present study, we have addressed the consequences of c-FLIP ablation in the ER stress response by investigating the effects of pharmacologically-induced ER stress in Wild Type (WT) and c-FLIP<sup>-/-</sup> MEFs. Surprisingly, c-FLIP<sup>-/-</sup> MEFs were found to be strikingly more resistant than WT MEFs to ER stress-mediated apoptosis. Analysis of Unfolded Protein Response (UPR) pathways revealed that Pancreatic ER Kinase (PERK) and Inositol-Requiring Enzyme 1 (IRE1) branch signalling is compromised in c-FLIP<sup>-/-</sup> cells when compared with WT cells. We found that c-FLIP modulates the PERK pathway by interfering with the activity of the serine threonine kinase AKT. Indeed, c-FLIP<sup>-/-</sup> MEFs display higher levels of active AKT than WT MEFs upon ER stress, while treatment with a specific AKT inhibitor of c-FLIP<sup>-/-</sup> MEFs subjected to ER stress restores the PERK but not the IRE1 pathway. Importantly, the AKT inhibitor or dominant negative AKT transfection sensitizes c-FLIP<sup>-/-</sup> cells to ER stress-induced cell death while the expression of a constitutively active AKT reduces WT cells sensitivity to ER stress-induced death. Thus, our results demonstrate that c-FLIP modulation of AKT activity is crucial in controlling PERK signalling and sensitivity to ER stress, and highlight c-FLIP as a novel molecular player in PERK and IRE1-mediated ER stress response.